



# Ocean

## A G R I G R O W

### Seaweed liquid extract

Guaranteed Analysis: N - P - K | 0.1 - 0.01 - 0.2

Derived from: *Ascophyllum nodosum*

Non Plant Food Ingredients: Preservatives (Citric Acid,  
Preservative mix)

See back label for precautionary  
statements and mixing instructions

[www.oceankl.com](http://www.oceankl.com)

Net Content:  
**10L (10.3kg)**



## COMPATIBILITY

It is recommended to carry out a test before preparing the mixture and to consult technically authorised personnel.

## MIXING INSTRUCTIONS

1. Empty and clean the spraying/mixing tank.
2. Add minimal volume of water
3. Mix well (or shake) the product before taking the required amount.
4. Start mixer and add this required amount of product – Keep mixing.
5. Top up the mixture with volume of water (or other foliar product) required for the surface to spray.
6. This product can be tank-mixed, under normal conditions of uses, with most of the commercial foliar treatment products including oligo-element concentrates and phytosanitary products.

Limit of other salt solubility and water electro-conductivity (EC) must be in agreement with appropriate chemical and agricultural rules.

## PRECAUTIONARY STATEMENTS

Keep out of reach of children.

Do not eat, drink or smoke when using this product. Store in a fresh, dry place and keep away from extreme temperatures.

Keep only in original container.

Use the product according to the instructions in this label. In case of doubt, please contact authorized technicians.

## HANDLING AND STORAGE

The product can separate in two phases, however this does not alter its efficiency. The product should be stored between 5° and 40 °C to provide optimum results. If risk of frost store above soil level. Avoid dramatic changes of temperatures. Oxidation of the extract and colour change from green to brown is normal and does not affect the efficiency of the product. Open bottles will oxidiz3 fast due to the contact with air.

Disclaimer: The recommendations and information provided comes from several trials and studies that have been undertaken, however, during the application of the product several factors out of our control may take place (tank mixes, application, climatic conditions, etc.). Any recommendation must serve as a guide and be adapted to local conditions.

CROPS	Dosage* L/Ha	Foliar application and periods of use
Both leafy/fruity vegetables	2-3 L/ha	1st (first) application 8 -10 days after transplanting and repeat during the crop cycle at 10 -14-day intervals.
Potato	2-3 L/ha	1st (first) initiation application of the tuber. 2nd (second) application 14 days after the first intervention.
Carrot	2-L/ha	1st (first) application as soon as the seedling reaches 15 cm in height and repeat the treatment every 14 days.
Artichoke	2-3 L/ha	1st (first) differentiation application of the central flower head and repeat every 7-10 days for 3-4 interventions.
Cereals	2 L/ha	In spring, apply at the beginning of the tillering.
Vine (table grapes)	2-3 L/ha	1st (first) before application of bunch distension (3-5 cm) 2nd (second) pre-flowering application with bunch still spreading (10 cm). Carry out 2-3 treatments after setting at intervals of 10 -14 days.
Vine (wine grapes)	2-3 L/ha	carry out 4-6 applications from budding to bunch ripening.
Olive tree	2-3 L/ha	Make 2-3 applications between the start of budding and flowering.
Drupaceous, citrus	2-3 L/ha	Make 3 applications: the 1st (first) at the beginning of flowering, the 2nd (second) at the fall of the petals and the 3rd (third) after the fruit sets.
Pomacee	2-3 L/ha	Make 3 applications: the 1st (first) at germination, the 2nd (second) at the formation of the flower bud and the 3rd (third) at the fall of the petals.
Strawberry	2-3 L/ha	Make 3 applications: the 1st (first) at germination, the 2nd (second) before flowering and the 3rd (third) with the setting of the first fruits
<b>Fertigation:</b> apply 6 - 10 L/ha, according to crop development, repeating every 10-14 days if necessary.		
Application dosage for turf: 10L - 20L per Ha		



## SHAKE WELL BEFORE USE

Batch Reference / Manufacture date:

Expiry: 14 months after Manufacture Date.  
Use within 21 days of opening.